

Name: \_\_\_\_\_

Panther ID: \_\_\_\_\_

Exam 1 - MAC2311 -

Spring 2015

**Important Rules:**

1. Unless otherwise mentioned, to receive full credit you **MUST SHOW ALL YOUR WORK**. Answers which are not supported by work might receive no credit.
2. Please turn your cell phone off at the beginning of the exam and place it in your bag, **NOT** in your pocket.
3. No electronic devices (cell phones, calculators of any kind, etc.) should be used at any time during the examination. Notes, texts or formula sheets should **NOT** be used either. Concentrate on your own exam. Do not look at your neighbor's paper or try to communicate with your neighbor.
4. Solutions should be concise and clearly written. Incomprehensible work is worthless.

1. (12 pts) These are True or False questions. No justification required. No partial credit. 2 points each.

(i) For all  $a > 0, b > 0$ ,  $\sqrt{a^2 + b^2} = a+b$  **True False**

(ii) The equation  $x^3 - 3x + 1 = 0$  has a solution in the interval  $[1, 2]$ . **True False**

(iii) If  $\lim_{x \rightarrow 3} f(x) = 1$  and  $\lim_{x \rightarrow 3} g(x) = -2$  then  $\lim_{x \rightarrow 3} [f(x) - 3g(x)] = 7$  **True False**

(iv) If  $x \neq 0$ ,  $\frac{\sin x}{x} = 1$  **True False**

(v) The function  $f(x) = \sec x$  is defined and is continuous for all real numbers  $x$ . **True False**

(vi) The inverse function of  $\sin x$  is  $\frac{1}{\sin x}$ . **True False**

2. (10 pts) Find all real solutions of the following equations (5 pts each):

(a)  $2 \cos x + 1 = 0$

(b)  $\log_{10}(x-15) - \log_{10} 2 = 3$

**3.** (12 pts) An object is dropped from the top of a building. Its position  $s(t)$  in feet above the ground  $t$  seconds after it was dropped is given by  $s(t) = 80 - 16t^2$ .

(a) (3 pts) When does the object hit the ground?

(b) (3 pts) Find the average velocity of the object during the first two seconds of its drop.

(c) (6 pts) Use limits to find the instantaneous velocity of the object at 2 seconds.

**4.** (12 pts) Sketch a graph of ONE function  $f(x)$  satisfying **all** of the following conditions.

(i) The function is defined and is continuous for all real  $x$ , except  $x = 0$  and  $x = 3$ ; the function is not defined at  $x = 0$  and  $x = 3$ .

(ii)  $\lim_{x \rightarrow 0^-} f(x) = +\infty$  and  $\lim_{x \rightarrow 0^+} f(x) = -\infty$ ;

(iii)  $\lim_{x \rightarrow 3^-} f(x) = 2$  and  $\lim_{x \rightarrow 3^+} f(x) = -2$ ;

(iv)  $\lim_{x \rightarrow -\infty} f(x) = 0$  and  $\lim_{x \rightarrow +\infty} f(x) = 2$ .

5. (30 pts) Find the following limits (5 pts each). If the limit is infinite or does not exist, specify so.

(a)  $\lim_{x \rightarrow 2} \frac{x^2 - 5x + 6}{4x - x^3}$

(b)  $\lim_{x \rightarrow +\infty} \frac{x^2 - 5x + 6}{4x - x^3}$

(c)  $\lim_{x \rightarrow 4} \frac{\sqrt{x} - 2}{x - 4}$

(d)  $\lim_{x \rightarrow 1} \frac{x^2 - 2x + 1}{|x - 1|}$

(d)  $\lim_{x \rightarrow 0} \frac{\sin(6x)}{x + \sin(2x)} =$

(e)  $\lim_{x \rightarrow 0^-} \cot x =$

6. (10 pts) Find, if possible, a value for the constant  $k$  which will make the function  $g(x)$  continuous everywhere. If you think there is no such  $k$ , justify why not.

$$g(x) = \begin{cases} \frac{\tan(kx)}{x} & \text{if } x < 0 \\ kx + 3 & \text{if } x \geq 0 \end{cases}$$

7. (a) (3 pts) Write the general  $(\epsilon, \delta)$  definition for  $\lim_{x \rightarrow a} f(x) = L$ .

Choose ONE of the parts (b) and (c). Only ONE will receive credit. Note the different point values.

(b) (7 pts) Use the  $(\epsilon, \delta)$  definition to prove  $\lim_{x \rightarrow 2} (2x-7) = -3$ .

(c) (12 pts) Use the  $(\epsilon, \delta)$  definition to prove  $\lim_{x \rightarrow 2} (2x^2-7) = 1$ .

**8.** (10 pts) Choose ONE of the following:

(a) State and prove the quadratic formula.

(b) Assuming the inequality  $\sin x \leq x \leq \tan x$  for  $x \in [0, \pi/2)$ , prove that  $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$ .