

Name: \_\_\_\_\_

Panther ID: \_\_\_\_\_

Exam 1 - MAC2311 - version 2 -

Summer B 2018

**Important Rules:**

1. Unless otherwise mentioned, to receive full credit you **MUST SHOW ALL YOUR WORK**. Answers which are not supported by work might receive no credit.
2. Please turn your cell phone off at the beginning of the exam and place it in your bag, **NOT** in your pocket.
3. No electronic devices (cell phones, calculators of any kind, etc.) should be used at any time during the examination. Notes, texts or formula sheets should **NOT** be used either. Concentrate on your own exam. Do not look at your neighbor's paper or try to communicate with your neighbor.
4. Solutions should be concise and clearly written. Incomprehensible work is worthless.

1. (12 pts) These are True or False questions. No justification required. No partial credit. 2 points each.

(i) If  $\lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^+} f(x)$  then  $f(x)$  is continuous at  $x = 3$ . **True False**

(ii) By subtraction,  $\lim_{x \rightarrow +\infty} (x^3 - 1000x^2) = \infty - \infty = 0$ . **True False**

(iii) For all  $a, b > 0$ ,  $\sqrt{a^2 + b^2} = a + b$  **True False**

(iv) The function  $f(x) = \frac{x-2}{\sqrt{x^2+1}}$  is defined and is continuous for all real numbers  $x$ . **True False**

(v) If  $y = L$  is a horizontal asymptote for the function  $y = f(x)$ , then it is possible for the graph of  $f$  to intersect the line  $y = L$  infinitely many times. **True False**

(vi)  $x = 5$  is a removable discontinuity for the function  $f(x) = \frac{x^2 - 25}{x - 5}$ . **True False**

2. (12 pts) An object is dropped from the top of a building. Its position  $s(t)$  in feet above the ground  $t$  seconds after it was dropped is given by  $s(t) = 400 - 16t^2$ .

(a) (2 pts) When does the object hit the ground?

(b) (4 pts) Find the average velocity of the object in the time interval  $0 \leq t \leq 2$  seconds.

(c) (6 pts) Use limits to find the instantaneous velocity of the object when  $t = 2$  seconds.

4. (30 pts) Find the following limits. If the limit is infinite or does not exist, specify so.

(a) (4 pts)  $\lim_{x \rightarrow 3} \frac{9x - x^3}{x^2 - x - 6}$

(b) (4 pts)  $\lim_{x \rightarrow 4} \frac{\sqrt{x} - 2}{4 - x}$

(c) (4 pts)  $\lim_{x \rightarrow 2} \frac{x}{x^2 - 4x + 4}$

(d) (4 pts)  $\lim_{x \rightarrow -\infty} \frac{4x^5 + 3x - 2}{3x^5 + 4}$

(e) (4 pts)  $\lim_{x \rightarrow +\infty} \sin\left(\frac{\pi x}{6x + 1}\right)$

(f) (5pts)  $\lim_{x \rightarrow 0} \frac{\tan^3(3x)}{x \sin^2(2x)}$

(g) (5pts)  $\lim_{x \rightarrow 0} x \sin(1/x)$

5. (9 pts) Given the function  $g(x) = \begin{cases} kx^2 + 2 & \text{if } x \leq 0 \\ \frac{\sin(kx)}{x} & \text{if } x > 0 \end{cases}$

is there a value of the constant  $k$  which will make  $g(x)$  continuous everywhere? Justify using limits.

6. (12 pts) Sketch the graph of ONE function  $f(x)$  satisfying ALL of the following conditions.

(i) The function is defined for all real numbers;

(ii) The function is continuous everywhere except  $x = 0$  and  $x = 3$ ;

$$(iii) \lim_{x \rightarrow 0^-} f(x) = -\infty, \quad f(0) = 4, \quad \lim_{x \rightarrow 0^+} f(x) = 4;$$

$$(iv) \lim_{x \rightarrow 3} f(x) = -\infty, \quad f(3) = 0, \quad ;$$

$$(v) \lim_{x \rightarrow -\infty} f(x) = 2, \quad \lim_{x \rightarrow +\infty} f(x) = 0.$$

7. (10 pts) Use the Intermediate Value Theorem to show that the equation  $x^4 = 1 - x^3$  has at least two distinct real solutions and locate each solution in an interval of length 0.5. Justify your work.

**8.** (10 pts) Choose ONE of the following. Only ONE will receive credit.

(A) State and prove the quadratic formula.

(B) Prove that  $\sin x \leq x \leq \tan x$  for any angle  $x \in [0, \pi/2)$ .