MAC 2233, Fall 2017

Exam #4

December 6, 2017

Name ____

- You will be told when to begin the work and when to terminate work on the examination. You must stop when instructed. Points may be deducted in case of violations.
- Please show your work to support your answers that require calculations. Correct but unsupported answers may not be given full credit.
- The use of a cell phone or other electronic communication devices during the examination is not allowed. The exam will be canceled and a grade of "0" will be assigned to anyone who opens a cell phone during the examination or if one is found on their seat or hand.

No calculators are allowed!

$$D(x,y) = f_{xx}(x,y) \cdot f_{yy}(x,y) - \left(f_{xy}(x,y)\right)^2$$

Honor Code: On my honor, I have neither received nor given any aid during this examination.

Signature:

1. (X points each) Find the second partial derivatives (mixed derivative included)

(a)
$$f(x,y) = e^{xy}$$

 $f_x = e^{xy}, y = y e^{xy}$
 $f_y = e^{xy}, x = x e^{xy}$

$$f_{xx} = ye^{xy}.y = y^2e^{xy}$$

$$f_{yy} = xe^{xy}.x = x^2e^{xy}$$

$$f_{xy} = |e^{xy}+y\cdot e^{xy}.x$$

$$= |e^{xy}+xye^{xy}|$$

$$= |e^{xy}(|+xy)|$$

$$f_{x} = \frac{2x}{x^{2} + y} = 2x(x^{2} + y)$$

$$f_{y} = \frac{1}{x^{2} + y} = (x^{2} + y)^{-1}$$

$$f_{x} = \frac{2x}{x^{2} + y} = 2x(x^{2} + y)$$

$$f_{xx} = \frac{2(x^{2} + y)^{2} + 2x(-1)(x^{2} + y)^{2}}{2(x^{2} + y)^{2} - (x^{2} + y)^{2}}$$

$$f_{y} = \frac{1}{x^{2} + y} = (x^{2} + y)^{2}$$

$$f_{yy} = -(x^{2} + y)^{2} \cdot 2x = -(x^{2} + y)^{2}$$

$$f_{yy} = -(x^{2} + y)^{2} \cdot 2x = -(x^{2} + y)^{2}$$

2. The demand function for peanut butter is

$$D_1(p_1, p_2) = 800 - 3p_1 - 4p_2$$

while that for a second commodity is

$$D_2(p_1, p_2) = 500 - 2p_2^3 - \frac{p_1}{2}$$

Is the second commodity more likely to be jelly or bread? Explain. [hint: are the two commodities substitute or complementary?]

$$\frac{\partial D_i}{\partial \rho_2} = 0 - 0 - 4 = -4 < 0$$
The commodities are complementary. Hence
$$\frac{\partial D_2}{\partial \rho_i} = 0 - 0 - \frac{1}{2} = -\frac{1}{2} < 0$$
The commodities are complementary. Hence the second comm. is probably bread.

3. A grocer's daily profit from the sale of two brands of cat food is

$$P(x,y) = 5x^2 + 2xy - 460x - 7y^2 + 480y - 1100$$

dollars, where x is the price per can of the first brand and y is the price per can of the second. Currently the first brand sells for \$2 per can and the second for \$5 per can.

(a) Use marginal analysis to estimate the change in daily profit that will result if the grocer raises the price of the **first** brand by \$1 per can but keeps the price of the second brand unchanged.

$$\frac{\partial P}{\partial x} = (0 \times + 2y - 460)$$

$$\frac{\partial P}{\partial x} (2,5) = 10.2 + 2.5 - 460 = 30 - 460 = -430$$
The daily profit decreases by \$430.

(b) Use marginal analysis to estimate the change in daily profit that will result if the grocer raises the price of the **second** brand by \$1 per can but keeps the price of the first brand unchanged.

$$\frac{\partial P}{\partial y} = 2 \times -14y + 480$$

$$\frac{\partial P}{\partial y} (2.5) = 2 \cdot 2 - 14 \cdot 5 + 480 = 4 - 70 + 480$$

$$= 414 \quad \text{The daily profit increases by}$$

(c) Increasing the price of which brand will yield higher daily profit?

4. Find the critical points of the given function and classify each as a relative maximum, a relative minimum, or a saddle point.

(a)
$$f(x,y) = xy$$

$$f_{x} = y$$

$$f_{y} = x$$

$$f_{y} = x$$

$$f_{xx} = 0$$

$$C_{y}(t \cdot p + s) \cdot \begin{cases} y = 0 \\ x = 0 \end{cases}$$

$$C_{y}(t \cdot p + s) \cdot \begin{cases} y = 0 \\ x = 0 \end{cases}$$

$$C_{y}(t \cdot p + s) \cdot \begin{cases} y = 0 \\ x = 0 \end{cases}$$

$$C_{y}(t \cdot p + s) \cdot \begin{cases} y = 0 \\ x = 0 \end{cases}$$

$$C_{y}(t \cdot p + s) \cdot \begin{cases} y = 0 \\ x = 0 \end{cases}$$

$$C_{y}(t \cdot p + s) \cdot \begin{cases} y = 0 \\ x = 0 \end{cases}$$

$$C_{y}(t \cdot p + s) \cdot \begin{cases} y = 0 \\ x = 0 \end{cases}$$

$$C_{y}(t \cdot p + s) \cdot \begin{cases} y = 0 \\ x = 0 \end{cases}$$

$$C_{y}(t \cdot p + s) \cdot \begin{cases} y = 0 \\ x = 0 \end{cases}$$

$$C_{y}(t \cdot p + s) \cdot \begin{cases} y = 0 \\ x = 0 \end{cases}$$

$$C_{y}(t \cdot p + s) \cdot \begin{cases} y = 0 \\ x = 0 \end{cases}$$

$$C_{y}(t \cdot p + s) \cdot \begin{cases} y = 0 \\ x = 0 \end{cases}$$

$$C_{y}(t \cdot p + s) \cdot \begin{cases} y = 0 \\ x = 0 \end{cases}$$

$$C_{y}(t \cdot p + s) \cdot \begin{cases} y = 0 \\ x = 0 \end{cases}$$

$$C_{y}(t \cdot p + s) \cdot \begin{cases} y = 0 \\ x = 0 \end{cases}$$

$$C_{y}(t \cdot p + s) \cdot \begin{cases} y = 0 \\ x = 0 \end{cases}$$

$$f_{xx} = 0$$

$$f_{yy} = 0$$

$$f_{xy} = 1$$

$$\int_{-\infty}^{\infty} \frac{D(o_1o) = -1 < 0}{(o_1o) \text{ is a}}$$

$$\int_{-\infty}^{\infty} \frac{dd}{dt} = \text{point.}$$

(b)
$$f(x,y) = x^3 - 12x + y^2 + 2y + 2y$$

$$f_{x} = 3x^{2} - 12$$

$$f_{y} = 2y + 2$$

$$f_{yy} = 2$$

$$f_{xy} = 0$$

$$\frac{(vi4. pts)}{(3(x^2.4)=0)} = 0 \longrightarrow x^2 = 4 \qquad (2i-1), (2i+1)$$

$$2(y+1)=0 \qquad x=\pm 2$$

$$y=-1$$

$$0 = 6x \cdot 2 - 0^2 = 12x$$

$$(2_{i}-1);$$

$$(2_{i}-1) = (2 \cdot 2 = 24 > 0)$$


$$(2_{i}-1) \text{ is a velative minimum.}$$

$$f_{xx}(2_{i}-1) = 6 \cdot 2 = 12 > 0$$

$$(-2_{i}-1); \quad 0(-2_{i}-1) = (2 \cdot (-2) = -24 < 0 \implies (-2_{i}-1) \text{ is a saddle point}$$

$$(-2_{1}^{-1})$$
: $(-2_{1}^{-1}) = (2 \cdot (-2) = -24 < 0 \rightarrow)$ (-2_{1}^{-1}) is a saddle point

- 5. (5 extra credit points) Find the critical points of the functions in problem 1 and classify each as a relative maximum, a relative minimum, or a saddle point.



- (b) $f(x,y) = \ln(x^2 + y)$ (vit. pts: $\frac{2x}{x^2 + y} = 0$ $\frac{1}{x^2 + y} = 0$ doesn't have a solution, i.e., $f_y \neq 0 = 0$ no critical points
- 6. (2 extra credit points each) Determine if the following statement is true or false.
 - (a) (true / false)
 - A function f(x,y) has a relative maximum at (a,b) if the sign of both first derivatives change from positive to negative about the point (a, b).
 - (b) (true/false)
 - If a function f(x,y) has all first and second partial derivatives $f_x, f_y, f_{xx}, f_{yy}, f_{xy}, f_{yx}$ then $f_{xy} = f_{yx}$.