STA 2122 Lab

What do I know already? (Do not look up the answers to these questions. The purpose is to assess your current level of knowledge on these topics.)

A. In your own words describe what a measure of variation (dispersion) reveals about a data set.

B. Why is the standard deviation generally preferred to the variance as a measure of dispersion (variation)?

C. What useful information can Chebyshev's theorem provide us with for a set of data? What assumptions about the shape of the distribution of the data does the theorem make?

D. In a left-skewed distribution, is the value of the mean larger or smaller than the median? What about in a right-skewed distribution?

Learning Objectives: (Click the learning objectives below for a short clip on the topic.)

Define and Calculate the Range (3) Determine when the Range is an Appropriate Measure of Dispersion (2) Define the Standard Deviation in Relation to Variance (1) Calculate the Sample Standard Deviation for a Set of Data (3) Compare the Usefulness of Variance and Standard Deviation Regarding the Units (2) State Chebyshev's Theorem (1) Calculate the Minimum Percentage of Data Inside a Symmetric Interval (3) Calculate the Maximum Percentage of Data Outside a Symmetric Interval (3) Create an Interval that Captures At Least 1-1/k^2 of the Data (4) Locate the Mean, Median, and Mode in Skewed and Symmetric Distributions (1) Categorize a Distribution as Either Left Skewed, Right Skewed, or Symmetric (2)

Exercises:

- In a left-skewed distribution, the median is 64. Which of the following values could be the mean for the distribution? (select all that apply)
 A. 62.1
 B. 61.3
 C. 65.4
 D. 67.0
 E. 64.2
 F. 62.9
- 2. Use the provided values {6, 2, 9, 1, -3, 5, 4} to find the following sum: $\sum_{i=1}^{7} (x_i 4)^2$
- 3. If the standard deviation for a sample of data is 1.3 ft, what is the variance for that sample of data?
- 4. The following frequency distribution is for grades in a course that has a maximum of 130 available points. Find the relative frequency for the class that includes grades between 70 and 80 points.

Frequency Distribution - Grades

	Grades		
lower		upper	frequency
20	<	30	1
30	<	40	0
40	<	50	13
50	<	60	26
60	<	70	53
70	<	80	54
80	<	90	31
90	<	100	15
100	<	110	5
110	<	120	1
120	<	130	1
			200

STA 2122 Lab

- In a right-skewed distribution, the median is 16.2. Which of the following values could be the mean for the distribution? (select all that apply)
 A. 12.3 B. 17.8 C. 15.4 D. 18.1 E. 16.2 F. 13.4
- 6. A manufacturer that makes microprocessors for cellular phones randomly selects four microprocessors from their production line in order to test them. These tests are done to determine the length of time the microprocessors function under extreme use. The manufacturer would like to calculate some measure of dispersion (variation) for the measurements taken from each sample of four microprocessors selected. Would the range be an acceptable option under these circumstances? Why or why not?
- 7. The NHTSA (National Highway Traffic Safety Administration) conducted a study of accidents involving motorcycles. The data included the ages of motorcyclists involved in accidents. If the researchers want to describe the typical age of a motorcyclist involved in a crash, which measure of the center is likely most appropriate?
 - A. Mean
 - B. Median
 - C. Mode
 - D. Range
 - E. Mean Absolute Deviation
- 8. Calculate the standard deviation for a set of 34 values that have the following summary values:

$$\sum_{i=1}^{n} x_i = 439 \text{ and } \sum_{i=1}^{n} x_i^2 = 7,223$$

- 9. According to a report by Common Sense Media, teens are spending an average of 9 hours per day in front of a screen for entertainment. The standard deviation for the time teens spend in front of a screen for entertainment is 2.2 hours. What is the minimum percentage of teens that spend between 5 and 13 hours in front of a screen for entertainment each day?
- 10. The number of friends each Facebook user has is an example of: (select all that apply)
 - A. continuous data
 - B. discrete data
 - C. qualitative data
 - D. nominal level data
 - E. quantitative data

STA 2122 Lab

- 11. A camera manufacturer has produced a camera shutter that has an average lifespan of 150,000 actuations with a standard deviation of 12,030 actuations. What can be said about the percentage of these shutters that will last for more than 185,000 actuations? (hint: use Chevyshev's Theorem)
- 12. Civil engineering graduate students weighed a random sample of vehicles passing over a small bridge here in Miami. The standard deviation for the weights of the sampled cars was 953 pounds. What unit of measurement would the variance of the data have?
 A. Pounds
 B. Root Pounds
 C. Pounds Squared
 D. Kilograms
- 13. The average weight of cars on U.S. roads is 4,009 pounds. If we assume the standard deviation for the weights of the cars is 919 pounds, create an interval that would capture the weights of at least 75% of all cars on U.S. roads.
- 14. Salary data was collected for a local university. The average salary for faculty at the university was \$73,661. The median salary was \$86,810, and the most common salary (the mode) for the collected data set was \$89,200. Is the distribution likely left skewed, right skewed, or symmetric?
- 15. According to a report by Common Sense Media, teens are spending an average of 9 hours per day in front of a screen for entertainment. The standard deviation for the time teens spend in front of a screen for entertainment is 2.2 hours. What is the maximum percentage of teens that spend less than 4 hours in front of a screen for entertainment each day?
- 16. The distribution below appears to be?



