
Student Number:

Exam Number:

Read Me First: *Show all essential work very neatly. Use correct notation when presenting your computations and arguments. Write using complete sentences. Be careful. Remember this: "=" denotes "equals" , " \Rightarrow " denotes "implies" , and " \Leftrightarrow " denotes "is equivalent to". Do not "box" your answers. Communicate. Show me all the magic on the page.*

1. (30 pts.) Very basic derivatives Provide the first derivative for each of the following functions.

(a) $f(x) = \log_{50}(x)$ $f'(x) =$

(b) $f(x) = 50^x$ $f'(x) =$

(c) $f(x) = \tan^{-1}(x)$ $f'(x) =$

(d) $f(x) = e^x$ $f'(x) =$

(e) $f(x) = \sin(x)$ $f'(x) =$

(f) $f(x) = \cos^{-1}(0)$ $f'(x) =$

(g) $f(x) = \tan(x)$ $f'(x) =$

(h) $f(x) = \sec(x)$ $f'(x) =$

(i) $f(x) = \csc(x)$ $f'(x) =$

(j) $f(x) = \cot(x)$ $f'(x) =$

(k) $f(x) = \sin^{-1}(x)$ $f'(x) =$

(l) $f(x) = \ln(x)$ $f'(x) =$

(m) $f(x) = \sec^{-1}(x)$ $f'(x) =$

(n) $f(x) = x^{50}$ $f'(x) =$

(o) $f(x) = \cos(x)$ $f'(x) =$

2. (30 pts.) Very basic antiderivatives Give each of the following antiderivatives. Do not forget the arbitrary constant.

(a) $\int \cos(x) \, dx =$

(b) $\int x^{50} \, dx =$

(c) $\int \sin(x) \, dx =$

(d) $\int \sec(x) \cdot \tan(x) \, dx =$

(e) $\int e^x \, dx =$

(f) $\int \frac{1}{1 + x^2} \, dx =$

(g) $\int \frac{1}{(1 - x^2)^{1/2}} \, dx =$

(h) $\int x^{-1} \, dx =$

(i) $\int \csc^2(x) \, dx =$

(j) $\int \frac{1}{|x| (x^2 - 1)^{1/2}} \, dx =$

(k) $\int \csc(x) \cdot \cot(x) \, dx =$

(l) $\int \sec^2(x) \, dx =$

(m) $\int \ln(50) \, dx =$

(n) $\int 50^x \, dx =$

(o) $\int 0 \, dx =$

3. (20 pts.) Some basic limits For each of the following, find the limit if it exists. If the limit does not exist, say so. Be as precise as possible in doing this. [Warning: At least one of these involves the definition of the derivative!!]

$$(a) \quad \lim_{x \rightarrow -7^+} \frac{x^2 - 49}{7x + 49} =$$

$$(b) \quad \lim_{x \rightarrow +\infty} \frac{4 - 9x^2 + 3x^5}{2x^5 - 11x + 11} =$$

$$(c) \quad \lim_{\theta \rightarrow 0} \frac{\sin(27\pi^3\theta)}{\tan(9\pi^2\theta)} =$$

$$(d) \quad \lim_{\theta \rightarrow 0} \frac{\tan((\pi/4) + \theta) - \tan(\pi/4)}{\theta} =$$

$$(e) \quad \lim_{x \rightarrow \infty} 2 \cdot \tan^{-1}(x) =$$

4. (20 pts.) More devious derivatives to do
Compute the first derivative of each of the following functions.
Do not simplify the algebra.

(a) $f(x) = 5 \cdot x^{1/5} - \sec(x) + 30 \cdot \sin^{-1}(x) + \tan(x^4) - 8 \cdot e^{\ln(\pi/4)}$

$$f'(x) =$$

(b) $y = \sec(x^3) \cdot e^{3x^2 - 4}$

$$\frac{dy}{dx} =$$

(c) $g(\theta) = \frac{\tan(\theta) - 4 \cdot e^{\sin(\theta)}}{\theta^2}$

$$\frac{dg(\theta)}{d\theta} =$$

(d) $h(t) = \log_{10}(3t) - 4^{10t} + (10 \cdot t^3)^2 + \ln(23) \cdot x$

$$h'(t) =$$

5. (25 pts.) More ignominious integrals Evaluate each of the following antiderivatives or indefinite integrals.

(a) $\int 6x^5 - \frac{9}{x} - 5 \cdot \cos(5x) \, dx =$


(b) $\int e^{9x+3} - \frac{52}{x^2+1} + \frac{20 \cdot x}{x^2+1} \, dx =$

(c) $\int 4 \cdot \cos^3(2\theta) \cdot \sin(2\theta) \, d\theta =$

(d) $\int (12t + 6) \cdot (t^2 + t) \, dt =$

(e) $\int \cos^3(x) \, dx =$

6. (10 pts.) Let $f(x) = x^4 - x^3$. Provide a complete analysis of this function on the back of Page 5. Then, below, plot the zero(s), the critical point(s), and the inflection point(s) of f . Finally, connect the dots and display the behavior of f at $\pm\infty$. Label very carefully.



7. (10 pts.) A child 5 ft. tall is walking at a rate of 4 ft./sec. toward a streetlight 20 feet high.

(a) At what rate is his shadow length changing?

(b) How fast is the tip of his shadow moving??

[Hints: (1) Draw a picture. (2) Look for similar triangles.]

8. (5 pts.) Give an equation for the line tangent to the graph of $f(x) = \sin(3x)$ at $x = -\pi/12$.

9. (10 pts.) Consider the equation $x^3 + 4x - 1 = 0$. Citing appropriate theorems, prove that this equation has exactly one solution on the real line. [Hints: You must show there is at least one solution. You must also show there is no more than one. Separate arguments are needed. Study $f(x) = x^3 + 4x - 1$.]

10. (10 pts.) Provide a complete evaluation each of the following limits. In particular, point out any use of L'Hopital's Rule, and when *squeezing*, give a complete argument.

(a)
$$\lim_{t \rightarrow \infty} \frac{32 + \sin(\pi \cdot t)}{t} =$$

(b)
$$\lim_{x \rightarrow 0} \frac{e^{3x} + e^{-3x} - 2}{2 - 2\cos(3x)} =$$

11. (10 pts.) A rectangular area of 100 square ft. is to be fenced off. Two opposite sides will use fencing costing \$1 per foot, and the remaining two sides require fencing costing \$2 per foot. Find the dimensions of the rectangle that requires the least cost. [You must provide a sufficiently complete argument to convince skeptics that your relative extremum is absolute!!]

12. (5 pts.) Solve the following initial value problem:
 $y'(x) = 4 \cdot \sec(x)$ and $y(\pi/3) = 8$.

13. (5 pts.) Using implicit differentiation, compute d^2y/dx^2 when $x^4 + y^2 = 1$.

14. (5 pts.) Use logarithmic differentiation to find dy/dx when

$$y = (5^{\tan(x)}) / (30^{\ln(27\cos(x))}).$$

15. (5 pts.) Give a complete $\epsilon - \delta$ proof that

$$\lim_{x \rightarrow 2} (10x - 5) = 15.$$