


---

**READ ME FIRST:** Show all essential work very neatly. Use correct notation when presenting your computations and arguments. Write using complete sentences. Be careful. Remember this: " $=$ " denotes "equals" , " $\Rightarrow$ " denotes "implies" , and " $\Leftrightarrow$ " denotes "is equivalent to". Since the answer really consists of all the magic transformations, do not "box" your final results. Communicate. Show me all the magic on the page.

---

1. (15 pts.) (a) Sketch very carefully the bounded region bounded by the curves  $y = -x^2$  and  $y = x - 2$  on the coordinate system provided. Label very carefully. (b) Simply write the definite integral,  $dx$ , that yields the area of the region. (c) Evaluate the definite integral of part (b) using the Fundamental Theorem of Calculus.

(a)



(b) Area =

(c) =

---

2. (5 pts.) Give the definition of the function  $\ln(x)$  in terms of a definite integral and give its domain and range. Label correctly. (**Hint:** Complete the sentence, " $\ln(x) = \dots$  .")

---

3. ( 10 pts.) Compute (a) the net distance, and (b) the total distance, traveled between time  $t = 0$  and time  $t = \pi$  by a particle moving with the velocity function  $v = \cos(t)$ .


(a) Net\_Distance =

(b) Total\_Distance =

---

4. ( 10 pts.) Obtain the arc length along the curve defined by the equation  $y = (4 - x^2)^{1/2}$  from  $x = -(2)^{1/2}$  to  $x = (3)^{1/2}$ .

5. (10 pts.) (a) Sketch the region in the 1st quadrant enclosed by the curves defined by  $y = \sin x$ ,  $y = 0$ , and  $x = \pi/2$ . Suppose the region is revolved around the line defined by  $x = \pi$ . (b) Using the method of cylindrical shells, write down the definite integral used to compute the volume of the solid of revolution formed. **Don't evaluate the integral.** (c) Using the method of slicing [disks/washers here], write down the definite integral used to compute the same volume as in part (b). **Don't evaluate the integral.**

<p>(a)</p> 	<p>(b)</p> <p><math>V =</math></p>   <hr/> <p>(c)</p> <p><math>V =</math></p>
--	--

6. (10 pts.) Consider the definite integral below. (a) Write down the sum,  $S_4$ , used to approximate the value of the integral below if Simpson's Rule is used with  $n = 4$ . **Do not attempt to evaluate the sum. Be very careful.** (b) Write down the sum,  $T_4$ , used to approximate the value of the integral below if Trapezoid Rule is used with  $n = 4$ . **Do not attempt to evaluate the sum. Be very careful.**

$$\int_1^2 x^{1/3} dx$$

(a)  $S_4 =$

(b)  $T_4 =$

---

7. (20 pts.) Evaluate each of the following antiderivatives or definite integrals. Give exact values for definite integrals.

[5 pts./part]

(a)

$$\int (2t+1)\cos(t) \, dt =$$

(b)

$$\int_0^{(\pi/3)^{1/2}} (4x)\sin(x^2) \, dx =$$

(c)

$$\int_1^e \frac{1}{x[1 + (\ln(x))^2]} \, dx =$$

(d)

$$\int (t+1)e^{2t} \, dt =$$

---

8. (10 pts.) Suppose that  $n \geq 2$  is a positive integer. Show in detail how to derive the following reduction formula:

$$\int \sin^n(x) \, dx = -\frac{\sin^{n-1}(x)\cos(x)}{n} + \frac{n-1}{n} \int \sin^{n-2}(x) \, dx$$

---

9. (10 pts.) Differentiate each of the following functions.

(a)  $f(x) = \sec^{-1}(x)$   $f'(x) =$

(b)  $f(x) = \cot^{-1}(x)$   $f'(x) =$

(c)  $f(x) = 20^x$   $f'(x) =$

(d)  $f(x) = \sin^{-1}(x)$   $f'(x) =$

(e)  $f(x) = \log_{\pi}(x)$   $f'(x) =$

---

**Silly 10 Point Bonus:** Consider the following equation:

$$\tan^{-1}\left(\frac{x}{(1-x^2)^{1/2}}\right) = \int_0^x \frac{1}{(1-t^2)^{1/2}} \, dt$$

Is this equation true for each  $x$  in the open interval  $(-1,1)$ ?? On the back of Page 4 of 5, either provide me proof that the equation is true for every  $x$  in  $(-1,1)$ , or show me that there is some number  $x_0$  in the interval where the equation fails to be true.