
READ ME FIRST: Show all essential work very neatly. Use correct notation when presenting your computations and arguments. Write using complete sentences. Remember this: "=" denotes "equals", " \Rightarrow " denotes "implies", and " \Leftrightarrow " denotes "is equivalent to". Since the answer consists of all the magic transformations, do not "box" your final results. Communicate. Show me all the magic on the page, for I do not read minds. Eschew obfuscation.

1. (5 pts.) Using a complete sentence and appropriate notation, state precisely the First Part of the Fundamental Theorem of Calculus. [This is sometimes called the *Evaluation Theorem*.]

2. (5 pts.) Evaluate the following using the 1st Part of the Fundamental Theorem of Calculus.

$$\int_{-\pi/4}^{\pi/4} 2 \cos(2x) \, dx =$$

3. (5 pts.) Express the following sum using sigma notation, but do not attempt to find its numerical value.

$$1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \frac{1}{11} + \frac{1}{13} =$$

4. (5 pts.) Obtain the upper limit of summation and rewrite the summand in order to make the following equation true. **Do not attempt to evaluate the sum.**

$$\sum_{k=10}^{45} k^4 = \sum_{j=1}$$

5. (5 pts.) Express the following limit as a definite integral. Do not attempt to evaluate the integral.

$$L = \lim_{\max \Delta x_k \rightarrow 0} \sum_{k=1}^n (2 \cos^2 x_k^*) \Delta x_k ; \quad a = -\pi/2, \quad b = \pi/2.$$

L =

6. (5 pts.) Using complete sentences and appropriate notation, state the Second Part of the Fundamental Theorem of Calculus.

7. (10 pts.) Find each of the following derivatives.

(a) $\frac{d}{dx} \left[\int_{\sqrt{3}}^x \tan^{-1}(t) \, dt \right] =$

(b) $\frac{d}{dx} \left[\int_1^{\ln(x)} e^t \, dt \right] =$

8. (5 pts.) (a) (3 pts.) Give the definition of the function $\ln(x)$ in terms of a definite integral and give its domain and range. Label correctly. (**Hint:** To start, complete the sentence, " $\ln(x) = \dots$.")

(b) (2 pts.) Given $\ln(a) = 3$ and $\ln(b) = -2$, evaluate the following integral.

$$\int_1^{ab^2} \frac{1}{t} \, dt =$$

9. (5 pts.) Write the solution to the following initial value problem in terms of a definite integral taken with respect to the variable \mathbf{t} , so the differential denoting the variable of integration is \mathbf{dt} . DO NOT ATTEMPT TO EVALUATE THE DEFINITE INTEGRAL IN t THAT YOU OBTAIN.

$$\frac{dy}{dx} = \sec^2(x) - \sin(x), \quad y(\pi/4) = 1.$$

$$y(x) =$$

10. (5 pts.) Evaluate the definite integral below by expressing it in terms of u correctly and then evaluating the resulting integral using a formula from geometry.

$$\int_{e^{-3}}^{e^3} \frac{\sqrt{9 - (\ln(x))^2}}{x} dx ; u = \ln(x)$$

$$\int_{e^{-3}}^{e^3} \frac{\sqrt{9 - (\ln(x))^2}}{x} dx =$$

11. (10 pts.) Write each of the following two sums in closed form.

$$(a) \sum_{k=1}^{50} \left(\frac{1}{k} - \frac{1}{k+1} \right) =$$

$$(b) \sum_{k=0}^{100} (-1)^{k+1} \left(\frac{1}{2^k} \right) =$$

12. (10 pts.) If the function f is continuous on $[a,b]$, then the *net signed area* A between $y = f(x)$ and the interval $[a,b]$ is defined by

$$A = \lim_{n \rightarrow \infty} \sum_{k=1}^n f(x_k^*) \Delta x.$$

Reveal all the details in computing the numerical value of the net signed area of $f(x) = x^3$ over the interval $[0,2]$ using only the definition above with

$$x_k^*$$

the right end point of each subinterval in the regular partition.

13. (10 pts.) A particle moves with a velocity of $v(t) = \cos(t)$ along an s-axis. Find the displacement and total distance traveled over the time interval $[0, 3\pi/2]$.

Displacement =

Total_Distance =

14. (5 pts.) Evaluate the following limit:

$$L = \lim_{x \rightarrow 0} (1 + 2x)^{\frac{1}{x}} =$$

15. (10 pts.) Let the function g be defined by the equation

$$g(x) = \int_0^x \cos^4(t) \, dt - 3x$$

for $x \in (-\pi/2, \pi/2)$. Then

(a) $g(0) =$

(b) $g'(0) =$

(c) $g''(0) =$

(d) Determine the open intervals where g is increasing or decreasing. Be specific.

(e) Determine the open intervals where g is concave up or concave down. Be specific.

Silly 10 Point Bonus: Obtain the exact numerical value of the limit L in Problem 5 on Page 1 of 4. [Say where your work is, for it won't fit here.]