

READ ME FIRST: Show all essential work very neatly. Use correct notation when presenting your computations and arguments. Write using complete sentences. Remember this: "=" denotes "equals", " \Rightarrow " denotes "implies", and " \Leftrightarrow " denotes "is equivalent to". Since the answer consists of all the magic transformations, do not "box" your final results. Communicate. Show me all the magic on the page, for I do not read minds. Eschew obfuscation.

1. (5 pts.) Evaluate the following using the 1st Part of the Fundamental Theorem of Calculus.

$$\int_1^4 \frac{1}{x\sqrt{x}} dx = (-2x^{-1/2}) \Big|_1^4 = \dots = 1$$

2. (5 pts.) Using a complete sentence and appropriate notation, state precisely the First Part of the Fundamental Theorem of Calculus. [This is sometimes called the *Evaluation Theorem*.]

If $f(x)$ is continuous on $[a,b]$ and $g(x)$ is any antiderivative of f on $[a,b]$, then

$$\int_a^b f(x) dx = g(b) - g(a).$$

3. (5 pts.) Express the following sum using sigma notation, but do not attempt to find its numerical value.

$$-\frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \frac{1}{11} + \frac{1}{13} = \sum_{k=1}^6 \frac{(-1)^k}{2k+1}$$

4. (5 pts.) Obtain the upper limit of summation and rewrite the summand in order to make the following equation true. **Do not attempt to evaluate the sum.**

$$\sum_{k=10}^{45} 3^{2k} = \sum_{j=0}^{35} 3^{2j+20}$$

since $j = k - 10$ is equivalent to $k = j + 10$.

5. (5 pts.) Express the following limit as a definite integral. Do not attempt to evaluate the integral.

$$L = \lim_{\max \Delta x_k \rightarrow 0} \sum_{k=1}^n (2 \cos^5 x_k^*) \Delta x_k ; \quad a = -2\pi, \quad b = \pi.$$

$$L = \int_{-2\pi}^{\pi} 2 \cos^5(t) dt$$

6. (5 pts.) Using complete sentences and appropriate notation, state the Second Part of the Fundamental Theorem of Calculus.

// Let $f(x)$ be a function that is continuous on an interval I , and suppose that a is any point in I . If the function g is defined on I by the formula

$$g(x) = \int_a^x f(t) dt,$$

for each x in I , then $g'(x) = f(x)$ for each x in I .//

7. (10 pts.) Find each of the following derivatives.

(a) $\frac{d}{dx} \left[\int_x^\pi 4 \sin(t^3) dt \right] = -4 \sin(x^3)$

(b) $\frac{d}{dx} \left[\int_1^{1/x} \sin t dt \right] = (-x^{-2}) \sin\left(\frac{1}{x}\right)$

8. (5 pts.) (a) (3 pts.) Give the definition of the function $\ln(x)$ in terms of a definite integral and give its domain and range. Label correctly. (**Hint:** To start, complete the sentence, " $\ln(x) = \dots$ ".)

$$\ln(x) = \int_1^x \frac{1}{t} dt$$

for $x > 0$. The domain of the natural log function is $(0, \infty)$, and its range is the whole real line, $(-\infty, \infty)$.

(b) (2 pts.) Given $\ln(a) = 8$ and $\ln(b) = -3$, evaluate the following integral.

$$\int_1^{ab^2} \frac{1}{t} dt = \ln(a) + 2 \ln(b) = 8 + (2)(-3) = 2$$

9. (5 pts.) Write the solution to the following initial value problem in terms of a definite integral taken with respect to the variable t , so the differential denoting the variable of integration is dt . DO NOT ATTEMPT TO EVALUATE THE DEFINITE INTEGRAL IN t THAT YOU OBTAIN.

$$\frac{dy}{dx} = 2e^{x^4}, \quad y(1) = \pi/4.$$

$$y(x) = \frac{\pi}{4} + \int_1^x 2e^{t^4} dt$$

10. (5 pts.) Evaluate the definite integral below by expressing it in terms of u correctly and then evaluating the resulting integral using a formula from geometry.

$$\int_{e^{-5}}^1 \frac{\sqrt{25 - (\ln(x))^2}}{x} dx ; u = \ln(x)$$

$$\int_{e^{-5}}^1 \frac{\sqrt{25 - (\ln(x))^2}}{x} dx = \int_{-5}^0 \sqrt{25 - u^2} du = \frac{25\pi}{4}$$

since $du = 1/x dx$. The right-most integral provides the area of one fourth of a circle with a radius of 5 centered at the origin.

11. (10 pts.) Write each of the following two sums in closed form.

$$(a) \sum_{k=1}^n \left(\frac{1}{k+2} - \frac{1}{k+3} \right) = \frac{1}{3} - \frac{1}{n+3}$$

$$(b) \sum_{k=0}^{100} \left(\frac{1}{3^k} \right) = \frac{3}{2} \left[1 - \left(\frac{1}{3} \right)^{101} \right]$$

12. (10 pts.) If the function f is continuous on $[a,b]$, then the *net signed area* A between $y = f(x)$ and the interval $[a,b]$ is defined by

$$A = \lim_{n \rightarrow \infty} \sum_{k=1}^n f(x_k^*) \Delta x.$$

Reveal all the details in computing the numerical value of the net signed area of $f(x) = 8x^3$ over the interval $[0,1]$ using only the definition above with

$$x_k^*$$

the right end point of each subinterval in the regular partition. *Do not use the Fundamental Theorem, Part 1.*

$$\begin{aligned} A &= \lim_{n \rightarrow \infty} \sum_{k=1}^n f(x_k^*) \Delta x = \lim_{n \rightarrow \infty} \sum_{k=1}^n 8 \left(\frac{k}{n} \right)^3 \frac{1}{n} \\ &= \lim_{n \rightarrow \infty} \left(\frac{8}{n^4} \right) \sum_{k=1}^n k^3 = \lim_{n \rightarrow \infty} \left(\frac{8n^2(n+1)^2}{4n^4} \right) \\ &= \lim_{n \rightarrow \infty} 2 \left(1 + \frac{1}{n} \right)^2 = 2 \end{aligned}$$

since $\Delta x = 1/n$ and $x_k = k/n$ for $k = 0, 1, \dots, n$ are the end points of the intervals of the general regular partition.

13. (10 pts.) A particle moves with a velocity of $v(t) = 2\cos(2t)$ along an s-axis. Find the displacement and total distance traveled over the time interval $[\pi/4, \pi/2]$.

$$\text{Displacement} = \int_{\pi/4}^{\pi/2} v(t) \, dt = \int_{\pi/4}^{\pi/2} 2 \cos(2t) \, dt = \dots = -1$$

$$\text{Total_Distance} = \int_{\pi/4}^{\pi/2} |v(t)| \, dt = \int_{\pi/4}^{\pi/2} |2 \cos(2t)| \, dt = \dots = 1$$

14. (5 pts.) Evaluate the following limit:

$$L = \lim_{x \rightarrow 0} (1 + x)^{1/(6x)} = e^{1/6}$$

15. (10 pts.) Let the function g be defined by the equation

$$g(x) = \int_0^x e^{t^2} \, dt - 4x$$

for $x \in (-\infty, \infty)$. Then since

$$g'(x) = e^{x^2} - 4 = e^{x^2} - e^{\ln(4)} \quad \text{and} \quad g''(x) = 2xe^{x^2},$$

$$(a) \quad g(0) = \int_0^0 e^{t^2} \, dt - 4(0) = 0$$

$$(b) \quad g'(0) = -3$$

$$(c) \quad g''(0) = 0$$

(d) Determine the open intervals where g is increasing or decreasing. Be specific.

From properties of the exponential and natural logarithmic functions, it follows that $g' > 0$ when $x^2 > \ln(4)$, and $g' < 0$ when $x^2 < \ln(4)$. Thus, g is increasing when

$$x < -\sqrt{\ln(4)} \quad \text{or} \quad \sqrt{\ln(4)} < x,$$

and g is decreasing when

$$-\sqrt{\ln(4)} < x < \sqrt{\ln(4)}.$$

(e) Determine the open intervals where g is concave up or concave down. Be specific. [This is easy.]

After examining the second derivative, it is easy to see that g is concave up when $x > 0$, and g is concave down when $x < 0$.

Silly 10 Point Bonus: Reveal the magic in evaluating the following limit:

$$\lim_{x \rightarrow 0^+} \frac{1}{x} \int_{-x}^x \frac{\sec(t)}{\pi^2 + t^4} \, dt$$

[Say where your work is, for it won't fit here.]

An extended discussion of this problem may be found in the TestTomb: Fall, 2005.