

**READ ME FIRST:** Show all essential work very neatly. Use correct notation when presenting your computations and arguments. Write using complete sentences. Remember this: "=" denotes "equals" , " $\Rightarrow$ " denotes "implies" , and " $\Leftrightarrow$ " denotes "is equivalent to". Since the answer consists of all the magic transformations, do not "box" your final results. Communicate. Show me all the magic on the page, for I do not read minds. Eschew obfuscation.

1. (5 pts.) Evaluate the following using the 1st Part of the Fundamental Theorem of Calculus.

$$\int_1^4 \frac{1}{x\sqrt{x}} dx =$$

2. (5 pts.) Using a complete sentence and appropriate notation, state precisely the First Part of the Fundamental Theorem of Calculus. [This is sometimes called the *Evaluation Theorem*.]

3. (5 pts.) Express the following sum using sigma notation, but do not attempt to find its numerical value.

$$-\frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \frac{1}{11} + \frac{1}{13} =$$

4. (5 pts.) Obtain the upper limit of summation and rewrite the summand in order to make the following equation true. **Do not attempt to evaluate the sum.**

$$\sum_{k=10}^{45} 3^{2k} = \sum_{j=0}$$

5. (5 pts.) Express the following limit as a definite integral. Do not attempt to evaluate the integral.

$$L = \lim_{\max \Delta x_k \rightarrow 0} \sum_{k=1}^n (2 \cos^5 x_k^*) \Delta x_k ; \quad a = -2\pi, \quad b = \pi.$$

$$L =$$

---

6. (5 pts.) Using complete sentences and appropriate notation, state the Second Part of the Fundamental Theorem of Calculus.

---

7. (10 pts.) Find each of the following derivatives.

(a)  $\frac{d}{dx} \left[ \int_x^\pi 4 \sin(t^3) dt \right] =$

(b)  $\frac{d}{dx} \left[ \int_1^{1/x} \sin t dt \right] =$

---

8. (5 pts.) (a) (3 pts.) Give the definition of the function  $\ln(x)$  in terms of a definite integral and give its domain and range. Label correctly. (**Hint:** To start, complete the sentence, " $\ln(x) = \dots$  .")

(b) (2 pts.) Given  $\ln(a) = 8$  and  $\ln(b) = -3$ , evaluate the following integral.

$$\int_1^{ab^2} \frac{1}{t} dt =$$

---

9. (5 pts.) Write the solution to the following initial value problem in terms of a definite integral taken with respect to the variable **t**, so the differential denoting the variable of integration is **dt**. DO NOT ATTEMPT TO EVALUATE THE DEFINITE INTEGRAL IN **t** THAT YOU OBTAIN.

$$\frac{dy}{dx} = 2e^{x^4}, \quad y(1) = \pi/4.$$

$$y(x) =$$

10. (5 pts.) Evaluate the definite integral below by expressing it in terms of  $u$  correctly and then evaluating the resulting integral using a formula from geometry.

$$\int_{e^{-5}}^1 \frac{\sqrt{25 - (\ln(x))^2}}{x} dx ; u = \ln(x)$$

$$\int_{e^{-5}}^1 \frac{\sqrt{25 - (\ln(x))^2}}{x} dx =$$

11. (10 pts.) Write each of the following two sums in closed form.

$$(a) \sum_{k=1}^n \left( \frac{1}{k+2} - \frac{1}{k+3} \right) =$$

$$(b) \sum_{k=0}^{100} \left( \frac{1}{3^k} \right) =$$

12. (10 pts.) If the function  $f$  is continuous on  $[a,b]$ , then the *net signed area*  $A$  between  $y = f(x)$  and the interval  $[a,b]$  is defined by

$$A = \lim_{n \rightarrow \infty} \sum_{k=1}^n f(x_k^*) \Delta x.$$

Reveal all the details in computing the numerical value of the net signed area of  $f(x) = 8x^3$  over the interval  $[0,1]$  using only the definition above with

$$x_k^*$$

the right end point of each subinterval in the regular partition. *Do not use the Fundamental Theorem, Part 1.*

13. (10 pts.) A particle moves with a velocity of  $v(t) = 2\cos(2t)$  along an s-axis. Find the displacement and total distance traveled over the time interval  $[\pi/4, \pi/2]$ .

Displacement =

Total\_Distance =

14. (5 pts.) Evaluate the following limit:

$$L = \lim_{x \rightarrow 0} (1 + x)^{1/(6x)} =$$

15. (10 pts.) Let the function  $g$  be defined by the equation

$$g(x) = \int_0^x e^{t^2} dt - 4x$$

for  $x \in (-\infty, \infty)$ . Then

(a)  $g(0) =$

(b)  $g'(0) =$

(c)  $g''(0) =$

(d) Determine the open intervals where  $g$  is increasing or decreasing. Be specific.

(e) Determine the open intervals where  $g$  is concave up or concave down. Be specific.

**Silly 10 Point Bonus:** Reveal the magic in evaluating the following limit:

$$\lim_{x \rightarrow 0^-} \frac{1}{x} \int_{-x}^x \frac{\sec(t)}{\pi^2 + t^4} dt$$

[Say where your work is, for it won't fit here.]