


**READ ME FIRST:** Show all essential work very neatly. Use correct notation when presenting your computations and arguments. Write using complete sentences. Be careful. Remember this: " $=$ " denotes "equals" , " $\Rightarrow$ " denotes "implies" , and " $\Leftrightarrow$ " denotes "is equivalent to". Since the answer really consists of all the magic transformations, do not "box" your final results. Communicate. Show me all the magic on the page.

1. (10 pts.) (a) Sketch the region in the first quadrant enclosed by the curves  $y = \cos^{-1}(x)$ ,  $x = 0$ , and  $y = 0$ . (b) Using the method of disks or washers, write a single definite integral whose numerical value is the volume of the solid obtained when the region is revolved around the x-axis. **Do not evaluate the integral.** (c) Using the method of cylindrical shells, write down a definite integral to compute the same volume as in part (b). **Do not evaluate the integral.**

(a)



(b)

$V =$

---

(c)

$V =$

2. (10 pts.) (a) Write down, but do not attempt to evaluate the definite integral that gives the arc length along the curve defined by the equation  $y = (1/2)x^2$  from  $x = -(3)^{1/2}$  to  $x = (3)^{1/2}$ .

$L =$

(b) Suppose a spring has a natural length of 1 foot, and a force of 10 pounds is needed to compress the spring to a length of 8 inches. Write down the definite integral that gives the work done in stretching this spring from its natural length to a total length of 15 inches but do not attempt to evaluate the integral.

Work =

3. ( 10 pts.) Consider the definite integral below. (a) Write down the sum,  $S_4$ , used to approximate the value of the integral below if Simpson's Rule is used with  $n = 4$ . **Do not attempt to evaluate the sum. Be very careful.** (b) Write down the sum,  $T_4$ , used to approximate the value of the integral below if Trapezoid Rule is used with  $n = 4$ . **Do not attempt to evaluate the sum. Be very careful.**

$$\int_{20}^{21} x^{1/4} dx$$

(a)  $S_4 =$

(b)  $T_4 =$

4. (10 pts.) (a) Using literal constants A, B, C, etc., write the form of the partial fraction decomposition for the proper fraction below. Do not attempt to obtain the actual numerical values of the constants A, B, C, etc.

$$\frac{2x^2+4}{x(x+1)(x^2+1)^3} =$$

(b) If one were to integrate the rational function in part (a), one probably would encounter the integral below. Reveal, in detail, how to evaluate this integral.

$$\int \frac{x}{(x^2+1)^3} dx =$$

5. (10 pts.) (a) In making the substitution  $u = \tan(x/2)$ , show all the details in obtaining  $\sin(x)$  and  $\cos(x)$  in terms of  $u$  by correctly completing the equations below. **[Hint: It helps to draw a right triangle to represent the equation  $u = \tan(x/2)$ .]** (b) Show how to obtain the differential  $dx$  in terms of  $du$  by solving for  $x$  in the equation  $u = \tan(x/2)$  and differentiating. (c) Using a sentence or two, explain what this substitution does to integrals of rational functions of sine and cosine.

(a) (4 pts.)  
 $\sin(x/2) =$

$\cos(x/2) =$

$\sin(x) = \sin(2(x/2)) =$

$\cos(x) = \cos(2(x/2)) =$

(b) (4 pts.)

$u = \tan(x/2)$  implies  $x =$  .

So  $dx =$  .

(c) (2 pt.)

---

6. (10 pts.) Here are five trivial trigonometric integrals to evaluate.  
 [2 pts./part]

(a)  $\int \tan(x) dx =$

(b)  $\int \sec(x) dx =$

(c)  $\int \sin^2(x) dx =$

(d)  $\int \csc(x) dx =$

(e)  $\int \cot(x) dx =$

7. (40 pts.) Evaluate each of the following antiderivatives or definite integrals. Give exact values for definite integrals.

[5 pts./part]

(a)  $\int_0^{(\pi/4)^{1/2}} (8x)\sin(x^2) dx =$

(b)  $\int 2x \tan^{-1}(x) dx =$

(c)  $\int \frac{\sin^2(t)}{\cos(t)} dt =$

(d)  $\int (8x)\sin(x) dx =$

7. (Continued) Evaluate each of the following antiderivatives or definite integrals. Give exact values for definite integrals.

(e)  $\int x^2 e^x dx =$

(f)  $\int \frac{1}{(x^2 + 1)^{1/2}} dx =$

(g)  $\int_0^1 \ln(x^2 + 1) dx =$

(h)  $\int (1 - t^2)^{1/2} dt =$

---

**Silly 10 Point Bonus:** The magnitude of error in using Simpson's Rule to approximate the definite integral of  $f(x)$  on an interval  $[a,b]$  with  $n$  subintervals may be estimated using

$$\left| \int_a^b f(x) dx - S_n \right| = |ES_n| \leq \frac{K_4(b-a)^5}{180n^4},$$

provided the fourth derivative of  $f$ ,  $f^{(4)}(x)$ , is continuous on  $[a,b]$ , and  $|f^{(4)}(x)| \leq K_4$  for each  $x$  in  $[a,b]$ . If Simpson's rule is used to approximate the definite integral giving the exact value of  $\ln(2)$ , reveal how to determine how big must  $n$  be to ensure you have 2 decimal place accuracy. [Say where your work is for there isn't room here. Remember  $n$  must be even.]