

---

1. (15 pts.) Construct the truth tables for each of the following compound propositions in the space provided:

(a)  $p \leftrightarrow q$         (b)  $\neg p$         (c)  $p \vee q$  :    See pages 3 through 7.

---

2. (10 pts.) Write the contrapositive and converse of the statement, "If the bacon will fly, then the air is greasy," and label unambiguously. Which is equivalent to the original statement?

Contrapositive:        If the air is not greasy, then the bacon won't fly.

Converse:                If the air is greasy, then the bacon will fly.

Only the contrapositive is equivalent to the original statement.

---

3. (15 pts.) Let  $F(x,y)$  be the statement "x can fool y". The universe of discourse is all people. Use quantifiers to express each of the following statements:

(a) Frodo can fool everyone.         $(\forall x)F(\text{Frodo},x)$

(b) Gandalf can fool no one.         $\neg(\exists x)F(\text{Gandalf},x)$  or  
equivalently  $(\forall x)\neg F(\text{Gandalf},x)$

(c) Someone can fool everyone.         $(\exists x)(\forall y)F(x,y)$

---

4. (10 pts.) Determine the truth value of each of the following statements if the universe of discourse of each variable is the set of real numbers,  $\mathbb{R}$ .

(a)  $(\exists x)(\forall y)(x \cdot y = 0)$  True        (b)  $(\forall x)(\exists y)(x \cdot y = 0)$  True

---

5. (15 pts.) Suppose  $A = \{\emptyset, 3, 4\}$  and  $B = \{\emptyset, \{\emptyset\}\}$ . Then

$A \cap B = \{\emptyset\}$

$A \times B = \{(\emptyset, \emptyset), (3, \emptyset), (4, \emptyset), (\emptyset, \{\emptyset\}), (3, \{\emptyset\}), (4, \{\emptyset\})\}$

$|P(A)| = 2^{|A|} = 2^3 = 8$

---

6. (10 pts.) What can you say about sets A and B if  $B = A \cap B$ ? Prove your assertion. If  $B = A \cap B$ , then  $B \subseteq A$ . Proof:  $x \in B = A \cap B$  implies  $x \in A$  and  $x \in B$  which implies  $x \in A$ . // It turns out that the converse is also true.

---

7. (15 pts.) Suppose that  $f: \mathbb{R} \rightarrow \mathbb{Z}$  is the function defined by the formula  $f(x) = \lfloor x \rfloor$ , and suppose that  $A = \{x \in \mathbb{R} \mid -1 < x \leq \pi\}$  and  $B = \{x \in \mathbb{R} \mid -2 < x \leq 3\}$ . Using appropriate notation, give each of the following.  $A - B = (3, \pi]$   $f(A) = \{-1, 0, 1, 2, 3\}$

$f^{-1}(\{1, 2\}) = f^{-1}(\{1\}) \cup f^{-1}(\{2\}) = [1, 2) \cup [2, 3) = [1, 3)$

---

8. (10 pts.) Suppose  $g: A \rightarrow B$  and  $f: B \rightarrow C$  are functions. Prove that if both f and g are surjective, then  $f \circ g: A \rightarrow C$  is surjective. Proof: [1.6, 19b]