

MAC 2313 (Calculus III)
Test 4 Review- Spring 2015

1. a) Let G be the solid defined by the inequalities: $\sqrt{x^2 + y^2} \leq z \leq 20 - x^2 - y^2$. Find the coordinates of the centroid of G . b) Find the mass and center of gravity of the solid G enclosed by the portion of the sphere $x^2 + y^2 + z^2 = 2$ on or above the plane $z = 1$ if the density is $\delta = \sqrt{x^2 + y^2 + z^2}$.

2. a) State the fundamental theorem of line integral. b) Let $F(x, y) = (2xy + x)\vec{i} + (x^2 + 2y)\vec{j}$. b1) Show that F is conservative. b2) Find a potential function φ for F . b3) Evaluate the line integral $\int_C (2xy + x) dx + (x^2 + 2y) dy$ along the curve C parametrized by $\vec{r}(t) = \sqrt{1+t}\vec{i} + \sin^{-1} t \vec{j}$, $0 \leq t \leq 1$.

3. a) Set up, but do not evaluate, two iterated integrals equal to the surface integral $\int_{\sigma} y^2 z dS$, where σ is the portion of the cylinder $x^2 + z^2 = 4$ in the first octant between the planes $y = 0$, $y = 6$, $x = z$, and $x = 2z$. b) Consider the parametric surface given by $\mathbf{r}(u, v) = u\vec{i} + u \cos v \vec{j} + u \sin v \vec{k}$ with $0 \leq u \leq 4$ and $0 \leq v \leq \pi$. i) Find the area S of σ . ii) Find the mass M of σ if its density is $\delta(x, y, z) = x^2 + y^2 + z^2$. iii) Evaluate the surface integral $\int_{\sigma} x\sqrt{z} dS$ where σ is the portion of the paraboloid $z = x^2 + y^2$ in the first octant between the planes $z = 0$ and $z = 4$. iv) a) Find an equation for the tangent plane to the parametric surface σ given by: $\vec{r}(u, v) = 4u \cos v \vec{i} + u^2 \vec{j} + 3u \sin v \vec{k}$, at the point P corresponding to $(u, v) = (1, \pi/2)$.

4. Let $F(x, y) = (x^3 y + 4e^{-2x})\vec{i} + (\frac{x^4}{4} + y^2)\vec{j}$. a) Show that F is conservative. b) Find a potential function φ for F . c) Evaluate the line integral $\int_C (x^3 y + 4e^{-2x}) dx + (\frac{x^4}{4} + y^2) dy$ along the curve C parametrized by $\vec{r}(t) = \cos^3 t \vec{i} + \sin^3 t \vec{j}$, $0 \leq t \leq \pi$.

5. a) Let $\mathbf{F}(x, y, z) = (x^2 - 2yx)\vec{i} + (3y^2 - 2yz)\vec{j} + (5z^2 - 2xz)\vec{k}$. Find $\text{div}\mathbf{F}$ and $\text{curl}\mathbf{F}$. Evaluate the line integral $\int_C \text{curl}\mathbf{F} \cdot d\mathbf{r}$, where C is the triangle with vertices $(0,0)$, $(0,2)$ and $(2,0)$.

6. Let C be the curve given by $x = t$, $y = 3t^2$, $z = 6t^3$, $0 \leq t \leq 1$, and evaluate $\int_C xyz^2 ds$. b) Evaluate the line integral along C given by $C : x = t$, $y = t^2$, $z = 3t^2$, $0 \leq t \leq 1$, $\int_C \sqrt{1 + 30x^2 + 10y} ds$. c) Evaluate $\int_C y dx + z dy - x dz$ along the helix $x = \cos(\pi t)$, $y = \sin(\pi t)$, $z = t$ from the point $(1,0,0)$ to $(-1,0,1)$. d) Find the mass of a thin wire shaped in the form of the curve $x = e^t \cos t$, $y = e^t \sin t$, $(0 \leq t \leq 1)$ if the density function δ is proportional to the distance to the origin.

7. a) Find parametric equations for the paraboloid $z = x^2 + y^2$ in terms of the parameters θ and ϕ , where (ρ, θ, ϕ) are spherical coordinates of a point on the surface. b) Find a parametric representation of the portion of the sphere $x^2 + y^2 + z^2 = 9$ on or above the plane $z = 2$ in terms of the parameters r and θ , where (r, θ, z) are the cylindrical coordinates of a point on the surface.

8. a) Let $\mathbf{F}(x, y, z) = \sqrt{x^2 + y^2} \vec{k}$. Find the flux of \mathbf{F} across σ , where σ is the portion of the cone $\vec{r}(u, v) = u \cos v \vec{i} + u \sin v \vec{j} + 2u \vec{k}$, with $0 \leq u \leq \sin v$, $0 \leq v \leq \pi$. b) Let $\mathbf{F}(x, y, z) = x\vec{i} + y\vec{j} + 2z\vec{k}$. Find the flux of \mathbf{F} across σ , where σ is the portion of the paraboloid below the plane $z = y$, oriented by downward unit normals.

9. Let $\mathbf{F}(x, y, z) = (x^3 - e^y)\vec{i} + (y^3 + \sin z)\vec{j} + (z^3 - xy)\vec{k}$. Use the Divergence Theorem to find the flux of \mathbf{F} across σ , where:

a) σ is the boundary of the solid G , bounded above by the sphere $z = \sqrt{4 - x^2 - y^2}$ and below by the xy -plane, with outward orientation.

b) σ is the boundary of the cylindrical solid enclosed by $x^2 + y^2 = 4$, $z = 0$ and $z = 1$ with outward orientation.

10. a) Use Green's theorem to evaluate the line integral $\int_C (4y + \cos(1 + e^{\sin x})) dx + (2x - \sec^2 y) dy$, where C is the circle $x^2 + y^2 = 9$ going from $(0,3)$ to $(0,3)$ counterclockwise.

11. Review the Fundamental Theorem of Line Integral, Green's Theorem and the Divergence Theorem.